

4.9 Holditch's Theorem

WORKSHEET

The question


A movable point P has cartesian coordinates (x, y) , where x and y are functions of t . The polar coordinates of P with respect to the origin O are r and θ . Starting with the expression

$$\frac{1}{2} \int r^2 d\theta$$

for the area swept out by OP , obtain the equivalent expression

$$\frac{1}{2} \int \left(x \frac{dy}{dt} - y \frac{dx}{dt} \right) dt. \quad (*)$$

The ends of a thin straight rod AB lie on a closed convex curve C . The point P on the rod is a fixed distance a from A and a fixed distance b from B . The angle between AB and the positive x direction is t . As A and B move anticlockwise round C , the angle t increases from 0 to 2π and P traces a closed convex curve $\mathcal{D}$ inside C , with the origin O lying inside $\mathcal{D}$, as shown in the diagram.



Let (x, y) be the coordinates of P . Write down the coordinates of A and B in terms of a, b, x, y and t .

The areas swept out by OA, OB and OP are denoted by $[A], [B]$ and $[P]$, respectively. Show, using $(*)$, that

$$[A] = [P] + \pi a^2 - af$$

where

$$f = \frac{1}{2} \int_0^{2\pi} \left(\left(x + \frac{dy}{dt} \right) \cos t + \left(y - \frac{dx}{dt} \right) \sin t \right) dt.$$

Obtain a corresponding expression for $[B]$ involving b . Hence show that the area between the curves $\mathcal{C}$ and $\mathcal{D}$ is πab .

Hints

Try the question on your own first. If you get stuck, read the hints *one at a time*, and go back to the question after each one. Hints marked *Pause and try* are the key steps.



Ask yourself Check the theorem in the simplest case. If $\mathcal{C}$ is a circle of radius R , what curve does P trace, and what is the area of the ring?

STAGE 1 Area from coordinates



Ask yourself With $x = r \cos \theta$ and $y = r \sin \theta$, all functions of t , compute $x\dot{y} - y\dot{x}$, where dots denote $\frac{d}{dt}$.

STAGE 3 The area swept by OA



Ask yourself Apply $(*)$ to A . Expand $x_A\dot{y}_A - y_A\dot{x}_A$ and sort the terms by powers of a .



Pause and try Now find $[B]$. (*Hint: B is obtained from the formula for A by replacing a with what?*)

STAGE 4 Eliminating f



Ask yourself Both A and B go once round the same curve $\mathcal{C}$. What does that tell you about $[A]$ and $[B]$?

Try this next

- The area of an ellipse.** Use $(*)$ with $x = p \cos t$, $y = q \sin t$ to show that an ellipse with semi-axes p and q has area πpq .
- The circle case.** Prove the intersecting chords theorem: if two chords of a circle meet at P , the products of the lengths of their two segments are equal. Deduce Holditch's theorem for a circle.
- The midpoint.** Show that $f = \pi(a - b)$ means that when P is the midpoint of the rod, $f = 0$. What is the ring's area then, in terms of the rod's length L ?
- A triangle.** **HARDER** Use $(*)$ to find the area of the triangle with vertices $(0, 0)$, (x_1, y_1) , (x_2, y_2) , traversing its boundary, and show it equals $\frac{1}{2}|x_1y_2 - x_2y_1|$. This is the start of the "shoelace formula" for the area of any polygon.

Further reading

- ▶ **Mark Levi**, *The Mathematical Mechanic* (Princeton University Press). Uses physical reasoning, with rods, wheels and springs, to prove mathematical results, in the spirit of this section.
- ▶ **H. M. Schey**, *Div, Grad, Curl, and All That* (Norton). An informal introduction to vector calculus, including Green's theorem.
- ▶ **Hamnet Holditch**, “Geometrical theorem”, *Quarterly Journal of Pure and Applied Mathematics* 2 (1858). The original note, a single short paragraph with none of the fine print.

The full guided solution, and what the question leads to at university, is in Section 4.9:
https://keithmcnulty.github.io/step_book_mcnulty/ch04/holditch.html