

6.3 A Polynomial for $\sqrt{2} + \sqrt{3} + \sqrt{5}$

WORKSHEET

The question

(i) Show that, if $(x - \sqrt{2})^2 = 3$, then $x^4 - 10x^2 + 1 = 0$.

Deduce that, if $f(x) = x^4 - 10x^2 + 1$, then $f(\sqrt{2} + \sqrt{3}) = 0$.

(ii) Find a polynomial g of degree 8 with integer coefficients such that $g(\sqrt{2} + \sqrt{3} + \sqrt{5}) = 0$. Write your answer in a form without brackets.

(iii) Let a , b and c be the three roots of $t^3 - 3t + 1 = 0$.

Find a polynomial h of degree 6 with integer coefficients such that $h(a + \sqrt{2}) = 0$, $h(b + \sqrt{2}) = 0$ and $h(c + \sqrt{2}) = 0$. Write your answer in a form without brackets.

(iv) Find a polynomial k with integer coefficients such that $k(\sqrt[3]{2} + \sqrt[3]{3}) = 0$. Write your answer in a form without brackets.

Hints

Try the question on your own first. If you get stuck, read the hints *one at a time*, and go back to the question after each one. Hints marked *Pause and try* are the key steps.

? **Ask yourself** To find a polynomial with $\sqrt{2}$ as a root, you square $x = \sqrt{2}$ to get rid of the square root. How might you get rid of two different square roots, one at a time?

STAGE 2 Three square roots

? **Ask yourself** Let $x = \sqrt{2} + \sqrt{3} + \sqrt{5}$, so $x - \sqrt{5} = \sqrt{2} + \sqrt{3}$ is a root of f . Write $f(x - \sqrt{5}) = 0$ in the form $P(x) + \sqrt{5}Q(x) = 0$. How do you then eliminate $\sqrt{5}$?

STAGE 3 A cubic and a square root

? **Ask yourself** If $x = a + \sqrt{2}$, then $x - \sqrt{2}$ is a root of $t^3 - 3t + 1$. Substitute, separate the terms with and without $\sqrt{2}$, and square.

STAGE 4 Two cube roots

🕒 **Pause and try** Let $x = \sqrt[3]{2} + \sqrt[3]{3}$. Cube both sides. Can you write the result in terms of x and a single cube root?

Try this next

- 1 A primitive element.** Let $y = \sqrt{2} + \sqrt{3}$. Show that $y^3 = 11\sqrt{2} + 9\sqrt{3}$, and deduce that $\sqrt{2} = \frac{1}{2}(y^3 - 9y)$. Hence every number $p + q\sqrt{2} + r\sqrt{3} + s\sqrt{6}$ can be written as a polynomial in y .
- 2 Mixed roots.** Find a polynomial of degree 6 with integer coefficients that has $\sqrt{2} + \sqrt[3]{3}$ as a root.
- 3 Irreducibility.** Show that $x^4 - 10x^2 + 1$ has no rational roots and cannot be written as a product of two quadratics with integer coefficients. (*Hint:* try $(x^2 + ax + b)(x^2 - ax + c)$.) Deduce that $\sqrt{2} + \sqrt{3}$ has degree 4.
- 4 Trisection.** Show that $\cos 3\theta = 4\cos^3 \theta - 3\cos \theta$, and deduce that $\cos 20^\circ$ is a root of $8x^3 - 6x - 1 = 0$. Assuming this cubic has no rational roots (check this), explain why a 60° angle cannot be trisected with ruler and compasses.

Further reading

- ▶ **Ian Stewart**, *Galois Theory* (CRC Press). Field extensions, the tower law and the classical construction problems, written for undergraduates.
- ▶ **Richard Courant & Herbert Robbins**, *What Is Mathematics?* (Oxford University Press). Chapter 3 proves the impossibility of the Greek problems with no prerequisites.
- ▶ **Robin Hartshorne**, *Geometry: Euclid and Beyond* (Springer). Ruler-and-compass constructions treated with modern algebra.

The full guided solution, and what the question leads to at university, is in Section 6.3:
https://keithmcnulty.github.io/step_book_mcnulty/ch06/field-extensions.html